

# Reconstruction of weighted graphs by their spectrum

Lorenz Halbeisen, University of California at Berkeley  
Norbert Hungerbühler, University of Alabama at Birmingham

## Abstract

It will be shown that for almost all weights one can reconstruct a weighted graph from its spectrum. This result is the opposite to the well-known theorem of Botti and Merris which states that reconstruction of non-weighted graphs is in general impossible since almost all (non-weighted) trees share their spectrum with another nonisomorphic tree.

## 1 Notations and introduction

A weighted graph  $G$  is a pair  $(A, M)$ , where  $A = (A_{ij})$  is a symmetric real  $n \times n$  matrix with  $A_{ii} = 0$ , called adjacency matrix, and where the mass matrix  $M = \text{diag}(m_1, \dots, m_n)$  is a real diagonal  $n \times n$  matrix. The valence matrix  $D$  of  $G$  is defined to be the diagonal  $n \times n$  matrix with

$$D_{ii} = \sum_{j=1}^n A_{ij} =: d_i.$$

If all masses  $m_i$  are equal to 1 and if  $A_{ij} \in \{0, 1\}$  for all  $i, j$ , then  $G$  is just a simple graph and  $D$  its vertex degree matrix. If the masses  $m_i$  are positive and all  $A_{ij} \geq 0$ , we consider  $G$  as a model for a molecule consisting of  $n$  atoms with weights  $m_i$  and with  $A_{ij}$  being the elasticity constant of the chemical binding between  $m_i$  and  $m_j$ : That is, if  $v_i(t)$  denotes the scalar deviation of  $m_i$  at time  $t$  from its normal position, we have for every  $i$

$$-m_i \ddot{v}_i = \sum_{j=1}^n A_{ij}(v_i - v_j) = v_i D_{ii} - \sum_{j=1}^n A_{ij} v_j.$$

Hence, an eigenvibration  $v_j(t) = u_j e^{i\omega t}$ ,  $j = 1, \dots, n$ , of the molecule satisfies (in matrix notation)

$$\omega^2 M u = D u - A u$$

where  $u$  is the vector  $(u_1, \dots, u_n)$ . In other words the negative squares  $-\omega^2 = \lambda$  of the eigenfrequencies of the molecule are the spectrum of the (generalized) eigenvalue problem

$$\det(A - D - \lambda M) = 0.$$

Alternatively, we could regard this as a discrete model of an inhomogeneous drum consisting of  $n$  vertices bearing weights  $m_i$  and with  $A_{ij}$  being the elasticity constant between  $m_i$  and  $m_j$ . Such a discrete model can for example arise from discretizing the corresponding continuous problem for a numerical treatment.

Let us have a short look at the case of simple graphs when all  $m_i$  and all nonzero  $A_{ij}$  equal 1. The adjacency spectrum of a simple graph  $G$ , i.e. the eigenvalue spectrum of the adjacency matrix  $A$ , is widely studied (see e.g. [2] as a main reference). Nonisomorphic graphs (i.e. graphs whose adjacency matrices are not permutation similar) affording the same (adjacency) characteristic polynomial are called cospectral. Schwenk showed in [16] that almost all trees are cospectral. On the other hand the operator  $L = L(G) := A - D$  is the so called Laplace or Kirchhoff operator of  $G$  (Laplace operator because it is the discrete analogue of the Laplace differential operator, and Kirchhoff operator since  $L$  first occurred in the famous Matrix-Tree Theorem of G. Kirchhoff). In how far the spectrum of  $L$  reflects the spectral properties of molecules is discussed in [3], [5] and [8]. The relation between a simple graph and its Laplace spectrum is studied e.g. in [6], [7], [15] and [13]. As a general reference for recent results on spectral graph theory see [4]. One of the most striking results is the Theorem of Botti and Merris (see [1]) which generalizes the results of Schwenk [16], McKay [12] and Turner [17]:

**Theorem 1 (Botti-Merris)** *Let  $t_n$  be the number of nonisomorphic trees on  $n$  vertices and  $s_n$  the number of such trees  $T$  for which there exists a nonisomorphic tree  $\tilde{T}$  such that the polynomial identities*

$$d_\chi(yA(T) + zD(T) - xI) \equiv d_\chi(yA(\tilde{T}) + zD(\tilde{T}) - xI)$$

*in the three variables  $x$ ,  $y$  and  $z$  hold, simultaneously, for every irreducible character  $\chi$  of the symmetric group  $S_n$ . Then  $\lim_{n \rightarrow \infty} s_n/t_n = 1$ .*

Here,  $I$  is the identity and  $d_\chi$  denotes the *immanent*

$$d_\chi(B) = \sum_{p \in S_n} \chi(p) \prod_{i=1}^n b_{ip(i)}, \quad (1)$$

where  $B = (b_{ij})$  is an  $n \times n$ -matrix (e.g. for  $\chi = \varepsilon$ , the alternating character,  $d_\chi = \det$ ).

Techniques which are based on Sunada’s Trace Theorem have recently allowed to generate isospectral simple graphs which are not necessarily trees, see [9].

The results in [1] and [9] seem to indicate that in general it is impossible to reconstruct the structure of a molecule from its spectrum. However, we will see below that the case of weighted graphs offers the possibility of a reconstruction.

We will always identify the vector  $m \in \mathbb{R}^n$  with the mass matrix  $M(m) = \text{diag}(m)$ . For given  $m \in \mathbb{R}^n$  and a countable set  $\mathcal{A} \subset \mathbb{R}$  we denote by  $\mathcal{G}_{\mathcal{A},M(m)}$  the set of weighted graphs  $G = (A, M(m))$  with  $A = (A_{ij})$ ,  $0 \leq A_{ij} \in \mathcal{A}$ , and we will say  $G$  is a graph over  $\mathcal{A}$  and  $M(m)$ .

In this paper we look at the following problem: Given  $m = (m_1, \dots, m_n) \in \mathbb{R}_+^n$ ,  $\mathcal{A} \subset \mathbb{R}$  countable (e.g.  $\mathcal{A} = \{0, 1\}$  in the simplest case) and the Laplace spectrum  $\{x \in \mathbb{R} : \det(L - xM(m)) = 0\}$  of a graph  $(A, M(m)) \in \mathcal{G}_{\mathcal{A},M(m)}$ . Can you then compute the adjacency matrix  $A$  from this information? The naive answer would be just to compare the spectrum of every possible graph with the given spectrum. But first, this only works for a finite set  $\mathcal{A}$ , second, the number of simple graphs on  $n$  vertices grows superexponentially in  $n$  and hence the method is not practicable, and third it does not answer the question for which set of mass matrices (depending on  $\mathcal{A}$ ) the map  $A \mapsto \{x \in \mathbb{R} : \det(L - xM(m)) = 0\}$  is injective. The aim of this paper is to discuss conditions on the mass matrix  $M(m)$  such that the answer to this question is affirmative and to describe reconstruction algorithms. In a first part we will discuss the case  $\mathcal{A} = \{0, 1\}$  with a very strong growth condition on the masses  $m_i$  which implies that reconstruction of the graph from its spectrum is possible, and in a second part we will consider a general countable set  $\mathcal{A}$  and an algebraic condition which shows that for almost all mass matrices (in a sense that will be made precise) reconstruction of the adjacency matrix  $A$  is possible.

The conditions we give for reconstructability of weighted graphs are sufficient, but certainly far from being necessary. Therefore — although it seems not to be realistic to apply our results directly to real molecules, since the masses of the atoms of a molecule might not satisfy the growth or the algebraic condition we use — the given reconstruction results at least show that reconstructability is a phenomenon that does occur for weighted graphs. So, whether in concrete situations reconstruction is possible or not may be a matter of a more detailed analysis adapted to the case at hand.

## 2 The Laplacian spectrum of weighted graphs

In this section all nonzero  $A_{ij}$  are assumed to be 1, i.e., we consider the case  $\mathcal{A} = \{0, 1\}$ , and we ask for a condition on the mass matrix  $M$  which allows to decide which

masses are linked in a graph whose Laplace spectrum is known.

**Theorem 2** *There exist mass matrices  $M_0 = \text{diag}(m_1, \dots, m_n)$  such that the following is true: Let  $G = (A, M)$  and  $\tilde{G} = (\tilde{A}, \tilde{M})$  be weighted graphs over  $\mathcal{A} = \{0, 1\}$  such that  $M$  and  $\tilde{M}$  are permutation similar to  $M_0$ , then*

$$\det(L(G) - xM) \equiv \det(L(\tilde{G}) - x\tilde{M}) \quad (2)$$

*holds if and only if  $G$  and  $\tilde{G}$  are isomorphic graphs, i.e.  $A = P\tilde{A}P^{-1}$  and  $M = P\tilde{M}P^{-1}$  holds for a permutation matrix  $P$ . A possible choice is  $m_i = n^{(2^i)}$ .*

**Remark:** The proof will be constructive and provide an “algorithm” to reconstruct the adjacency matrix  $A$  from the roots of the polynomial  $\det(L - xM)$ .

The proof of Theorem 2 is based upon the following two elementary lemmas:

**Lemma 1** *Let  $q_1, \dots, q_n$  be a sequence of real numbers of at least geometric growth with constant  $s > 1$ , i.e.  $q_i \geq sq_{i-1}$  for  $i = 2, \dots, n$ , and  $q_1 > 0$ . Then*

$$\sum_{i=1}^n \delta_i q_i = \sum_{i=1}^n \tilde{\delta}_i q_i \quad (3)$$

*implies  $\delta_i = \tilde{\delta}_i$  for  $i = 1, \dots, n$ , provided that all  $\delta_i \in \{0, 1, \dots, \lfloor s - 1 \rfloor\}$ .*

**Proof**

We proceed by induction: For  $n = 1$  the assertion is trivial. On the other hand, using (3) we have for  $n > 1$

$$\begin{aligned} (\delta_n - \tilde{\delta}_n)q_n &= \sum_{i=1}^{n-1} (\tilde{\delta}_i - \delta_i)q_i & (4) \\ &\leq (s - 1) \sum_{i=1}^{n-1} q_i \\ &\leq (s - 1) \sum_{i=1}^{n-1} q_n \frac{1}{s^{n-i}} \\ &= q_n \left(1 - \frac{1}{s^{n-1}}\right) \\ &= q_n \varepsilon \end{aligned}$$

for an  $\varepsilon < 1$ . We may assume that  $\delta_n \geq \tilde{\delta}_n$  and hence we obtain from (4)

$$0 \leq \delta_n - \tilde{\delta}_n < 1.$$

Thus  $\delta_n = \tilde{\delta}_n$  and the assertion follows by induction.  $\square$

The second lemma we need in the proof of Theorem 2 is the following

**Lemma 2** *Let  $\mu_i = \nu^{(2^i)}$  for some  $\nu > 0$  and for  $i = 1, \dots, n$ . Then the set of the numbers*

$$q_{ij} = \frac{1}{\mu_i \mu_j} \prod_{k=1}^n \mu_k$$

*with  $i \neq j$  rearranged as a growing sequence has at least geometric growth with constant  $\nu$ .*

**Proof**

Consider  $a = \frac{q_{ij}}{q_{lm}}$  for  $\{i, j\} \neq \{l, m\}$ . We have  $a = \frac{\mu_l \mu_m}{\mu_i \mu_j} = \nu^{2^l + 2^m - 2^i - 2^j}$  and therefore the proof is complete if we can show that the exponent  $2^l + 2^m - 2^i - 2^j \neq 0$ . But this follows from Lemma 1 since  $2^l + 2^m = 2^i + 2^j$  would imply  $\{l, m\} = \{i, j\}$  which contradicts the assumption.  $\square$

Now we give the proof of Theorem 2:

**Proof**

We may assume that the vertex sets of  $G$  and  $\tilde{G}$  are already renumbered in such a way that  $M = \tilde{M} = M_0$ . Using (1) we easily find the following expansion

$$\begin{aligned} \det(L - xM) &= \begin{vmatrix} -d_1 - xm_1 & A_{12} & A_{13} & \dots & A_{1n} \\ A_{21} & -d_2 - xm_2 & A_{23} & \dots & A_{2n} \\ \vdots & \vdots & \ddots & & \vdots \\ A_{n1} & A_{n2} & \dots & \dots & -d_n - xm_n \end{vmatrix} = \quad (5) \\ &= (-1)^n x^n \prod_{i=1}^n m_i + (-1)^n x^{n-1} \sum_{i=1}^n d_i \prod_{j \neq i} m_j + \\ &\quad + (-1)^n x^{n-2} \sum_{i < j} (d_i d_j - A_{ij}^2) \prod_{k \notin \{i, j\}} m_k + \dots + \det(L). \end{aligned}$$

Now we use expansion (5) in the identity (2). Comparing the coefficients of  $x^{n-1}$  on both sides we conclude by Lemma 1 that

$$d_i = \tilde{d}_i \quad (i = 1, \dots, n) \quad (6)$$

since by our assumption on the masses  $m_i$  the ordered set of numbers  $q_i = \prod_{j \neq i} m_j$  is at least of geometric growth with constant  $n$  and  $d_i \in \{0, 1, \dots, n-1\}$ . Notice that by the theorem of Botti and Merris this cannot yet imply that the graphs  $G$  and  $\tilde{G}$  are isomorphic.

Comparing the coefficients of  $x^{n-2}$  and using (6) we obtain

$$\sum_{i < j} A_{ij} \prod_{k \notin \{i,j\}} m_k = \sum_{i < j} \tilde{A}_{ij} \prod_{k \notin \{i,j\}} m_k.$$

The numbers  $q_{ij} = \prod_{k \notin \{i,j\}} m_k$  obviously satisfy the hypothesis of Lemma 2 with  $\nu = n$  and hence we conclude (by applying Lemma 1 once more with  $s = 2$ ) that  $A_{ij} = \tilde{A}_{ij}$  and the proof is complete.  $\square$

Up to now, we consider two graphs as isospectral if they share the polynomial  $\det(L - xM)$ , i.e. the eigenvalues of both graphs coincide *counted with multiplicity*. Now we will show that even if we only require that two graphs have the same spectrum *as sets* they are isomorphic.

Let us consider a connected, weighted graph  $G$  with masses  $m_i = n^{(2^i)}$  as in Theorem 2. Then the following Proposition claims that the eigenvalues of the Laplacian spectrum of  $G$  are simple.

**Proposition 1** *Suppose  $\mathcal{A} = \{0, 1\}$  and let  $G \in \mathcal{G}_{\mathcal{A}, M(m)}$  be a connected weighted graph with masses  $m_i = n^{(2^i)}$ ,  $i = 1, \dots, n$ . Then the roots of the characteristic polynomial  $\det(L - xM(m))$  are simple.*

**Proof**

Let  $p(x) = (-1)^n \det(L - xM) = a_n x^n + \dots + a_2 x^2 + a_1 x + a_0$ . Since all roots  $\lambda_i = -\omega_i^2$  of  $p$  are negative real numbers, we have

$$a_k \geq 0 \quad \text{for } k = 0, \dots, n. \tag{7}$$

From (1) we get for  $k = 0, \dots, n$

$$a_k = \sum_{|I|=k} \det(L_I) \prod_{j \in I} m_j \tag{8}$$

where the sum is taken over all ordered subsets  $I$  of  $\{1, \dots, n\}$  of cardinality  $k$  and where  $L_I$  denotes the matrix obtained from  $L$  by deleting all rows and columns having a number in  $I$ . Of course, since the sum of the rows in  $L$  is zero,  $a_0 = \det(L) = 0$ .

Now, observe first that

$$|\det(L_I)| \leq n^{n-k} \tag{9}$$

for  $|I| = k$ . This follows from the fact that every column of  $L_I$  represents a vector of length at most  $n$ . On the other hand for  $1 \leq |I| < n$  we have

$$1 \leq |\det(L_I)| \tag{10}$$

since the graphs under consideration are assumed to be connected which implies that the matrices  $L_I$  are strongly diagonal dominated.

For simplicity we assume  $m_i = n^{(3^i)}$  in the proceeding of the proof. The arguments in case  $m_i = n^{(2^i)}$  are similar but more terms have to be taken into consideration. In order to obtain an estimate for the coefficients  $a_k$  we proceed as follows: The largest term in  $\sum_{|I|=k} \prod_{j \in I} m_j$  is obviously  $\Gamma_k := \prod_{j=n-k+1}^n m_j$ . All other terms are smaller or equal to  $\gamma_k := m_{n-k} \prod_{j=n-k+2}^n m_j$ . The quotient is  $\frac{\Gamma_k}{\gamma_k} = m_{n-k}^2$ . Since the total number of terms in the sum is  $\binom{n}{k}$  we obtain from (9), (10) and (8)

$$\left(1 - n^{n-k} \binom{n}{k} \frac{1}{m_{n-k}^2}\right) \prod_{j=n-k+1}^n m_j < |a_k| < \left(1 + \binom{n}{k} \frac{1}{m_{n-k}^2}\right) n^{n-k} \prod_{j=n-k+1}^n m_j. \quad (11)$$

An elementary calculation shows that for  $n \geq 1$  and  $k = 1, \dots, n$

$$n^{n-k} \binom{n}{k} \frac{1}{m_{n-k}^2} \leq \frac{1}{n^2}. \quad (12)$$

Inserting (12) in (11) yields

$$\left(1 - \frac{1}{n^2}\right) \prod_{j=n-k+1}^n m_j < |a_k| < \left(n^{n-k} + \frac{1}{n^2}\right) \prod_{j=n-k+1}^n m_j. \quad (13)$$

Using (13) we obtain that for  $k = 2, \dots, n-1$

$$a_k^2 - 4a_{k-1}a_{k+1} > 0 \quad (14)$$

provided that  $n \geq 3$  (the case  $n = 2$  is easily handled separately). Now the claim follows from the criterion of Kurtz on distinct roots of polynomials (see [11]).  $\square$

Combining Theorem 2 and Proposition 1 we obtain as a corollary

**Theorem 3** *There exist mass matrices  $M_0$  such that the following is true: Let  $G = (A, M)$  and  $\tilde{G} = (\tilde{A}, \tilde{M})$  be connected graphs over  $\mathcal{A} = \{0, 1\}$  such that  $M$  and  $\tilde{M}$  are permutation similar to  $M_0$ . Then  $G$  and  $\tilde{G}$  are isomorphic if and only if the Laplacian spectrum of  $G$  and  $\tilde{G}$  coincide as sets. A possible choice is  $m_i = n^{(2^i)}$ , where  $n$  is the number of masses.*

### Proof

According to Proposition 1 the Laplacian spectrum of both graphs consists of simple eigenvalues. Hence, by Viëta's Theorem, we conclude that  $\det(L(G) - xM) \equiv \mu \det(L(\tilde{G}) - x\tilde{M})$  for some  $\mu \neq 0$ . On the other hand the coefficient of  $x^n$  is

$(-1)^n \prod_{i=1}^n m_i$  in  $\det(L(G) - xM)$  and  $(-1)^n \prod_{i=1}^n \tilde{m}_i$  in  $\det(L(\tilde{G}) - x\tilde{M})$ , which for both cases is the same number since the mass matrices of  $G$  and  $\tilde{G}$  are permutation similar. Hence  $\mu = 1$  and the assertion follows from Theorem 2.  $\square$

**Algorithmic remark:** If one starts from the spectrum, the reconstruction algorithm works as follows. First, compute the polynomial  $\mu \det(L(G) - xM)$  by Viëta's Theorem and normalize it such that the coefficient of  $x^n$  is  $(-1)^n \prod_{i=1}^n m_i$ . Then find the valence matrix  $D$  using the coefficient of  $x^{n-1}$  as described in the proof of Theorem 2. Finally use this to compute the adjacency matrix  $A$  from the coefficient of  $x^{n-2}$ . (Notice that the proof of Lemma 1 can be used to determine recursively the values of the  $\delta_i$  from the value of the sum  $\sum_{i=1}^n \delta_i q_i$ .)

Inspection of the proof of Theorem 3 shows that the set of masses  $m \in \mathbb{R}^n$ , for which reconstruction is possible, contains a large open set (all sequences  $m_i$  which grow "fast enough"). However, the set of masses for which reconstruction is possible has also a part with a fine algebraic structure, as we will see in the next section. There, we consider a general countable set  $\mathcal{A}$  of possible values of elasticity constants and impose algebraic conditions on the masses to show that for almost all mass matrices the weighted adjacency matrix of a graph is determined by its Laplace spectrum.

### 3 $p$ -independent reals

In order to simplify the formulas, we use a multi-index notation: For  $i = (i_1, \dots, i_n) \in \mathbb{N}_0^n$  we write  $|i| := \max(i_1, \dots, i_n)$  and for  $m = (m_1, \dots, m_n) \in \mathbb{R}^n$  we define  $m^i := \prod_{k=1}^n m_k^{i_k}$ .

Let  $Q$  be a set of real numbers and  $p \in \mathbb{N}_0$ . We say that  $m \in \mathbb{R}^n$  is  $p$ -independent over  $Q$  if the following implication holds:

$$\sum_{i \in \mathbb{N}_0^n, |i| \leq p} q_i m^i = 0 \text{ and } q_i \in Q \text{ for all } i \in \mathbb{N}_0^n, |i| \leq p \implies q_i = 0 \text{ for all } i \in \mathbb{N}_0^n, |i| \leq p. \quad (15)$$

Notice that the set  $\{m_1, \dots, m_n\} \subseteq \mathbb{R}$  is algebraically independent over  $Q$  if and only if  $m = (m_1, \dots, m_n) \in \mathbb{R}^n$  is  $p$ -independent over  $Q$  for every  $p \in \mathbb{N}_0$ . So, the notion of  $p$ -independence is weaker than the notion of algebraic independence. For example,  $\sqrt[3]{2}$  and  $\sqrt[3]{5}$  are 2-independent but not algebraically independent over  $\mathbb{Q}$ .

**Lemma 3** *If  $Q \subset \mathbb{R}$  is countable and  $p \in \mathbb{N}_0$ , then the set  $\{m \in \mathbb{R}^n : m \text{ not } p\text{-independent over } Q\}$  is a meager and Lebesgue measure zero set in  $\mathbb{R}^n$ .*



**Proof**

For a fixed  $m = (m_1, \dots, m_{n-1}) \in \mathbb{R}^{n-1}$  let  $F_m$  be the set of all not identically vanishing polynomials  $f(x)$  with coefficients in  $Q \cup \{m_1, \dots, m_{n-1}\}$  of degree at most  $p$ . Because the set  $Q \cup \{m_1, \dots, m_{n-1}\}$  is countable and  $p$  is finite, the set

$$N(m) := \{x \in \mathbb{R} : f \in F_m \wedge f(x) = 0\}$$

is countable. Hence, for every  $m \in \mathbb{R}^{n-1}$ , the set  $N(m)$  is a meager and Lebesgue measure zero set in  $\mathbb{R}$  and by the theorems of Kuratowski–Ulam and Fubini (see, e.g., [14] or [10]) we get that the set  $\{m \in \mathbb{R}^n : m \text{ not } p\text{-independent over } Q\}$  is a meager and Lebesgue measure zero set in  $\mathbb{R}^n$ .  $\square$

Remember that there exist meager sets which do not have Lebesgue measure zero and vice versa. Moreover, one can cover the real line with a meager set and a set of Lebesgue measure zero.

## 4 The Reconstruction Theorem

In this section let  $C$  denote an arbitrary but fixed set of countably many real numbers. Then  $\mathcal{A} = \mathbb{Q}[C]$ , the smallest field containing  $C$  and the rational numbers, is countable as well. We show that if the set of masses  $m \in \mathbb{R}_+^n$  fulfills a suitable algebraic condition with respect to the set  $\mathcal{A}$ , then the adjacency matrix  $A(G)$  of a graph in  $\mathcal{G}_{\mathcal{A}, M(m)}$  is determined by its Laplace spectrum  $\{x \in \mathbb{R} : \det(L(G) - xM(m)) = 0\}$ . In particular, we will see that the set of masses  $m \in \mathbb{R}_+^n$  for which reconstruction is *not* possible is a meager and Lebesgue measure zero set in  $\mathbb{R}^n$ . Remember that for  $m \in \mathbb{R}_+^n$ ,  $M(m) = \text{diag}(m)$ , and that  $\mathcal{G}_{\mathcal{A}, M(m)}$  is the set of all weighted graphs  $G = (A, M(m))$  with  $A = (A_{ij})$  and  $0 \leq A_{ij} \in \mathcal{A}$ .

**Theorem 4** *Let  $m \in \mathbb{R}_+^n$  be 1-independent over  $\mathbb{Q}[C]$ , and let  $G = (A, M)$  and  $\tilde{G} = (\tilde{A}, \tilde{M})$  be graphs over  $\mathcal{A} = \mathbb{Q}[C]$  of order  $n$  such that  $M$  and  $\tilde{M}$  are permutation similar to  $M(m)$ . Then  $G$  and  $\tilde{G}$  are isomorphic if and only if their characteristic polynomials coincide, i.e., if  $\det(L(G) - xM) \equiv \det(L(\tilde{G}) - x\tilde{M})$ .*

**Proof**

It is easy to see that if  $G$  and  $\tilde{G}$  are isomorphic, then their characteristic polynomials coincide.

For the opposite implication we may assume that the vertex sets of  $G$  and  $\tilde{G}$  are already renumbered such that  $M = \tilde{M} = M(m)$ . We recall that  $p(x) = (-1)^n \det(L(G) - xM) = a_n x^n + \dots + a_2 x^2 + a_1 x + a_0$  with  $a_0 = 0$  and with  $a_{n-1} = \sum_{i=1}^n d_i \prod_{j \neq i} m_j$ .

If  $p(x)$  and  $\tilde{p}(x)$  coincide we have in particular  $\sum_{i=1}^n d_i \prod_{j \neq i} m_j = \sum_{i=1}^n \tilde{d}_i \prod_{j \neq i} m_j$ , hence  $\sum_{i=1}^n (d_i - \tilde{d}_i) \prod_{j \neq i} m_j = 0$ , and because  $m$  is 1-independent over  $\mathcal{A}$ , we have  $d_i = \tilde{d}_i$  for  $1 \leq i \leq n$ . Thus, the valence matrix is determined by the coefficient  $a_{n-1}$ . Comparing the coefficient  $a_{n-2} = \sum_{i < j} (d_i d_j - A_{ij}^2) \prod_{k \notin \{i, j\}} m_k$  of  $x^{n-2}$ , and using again that  $m$  is 1-independent over  $\mathcal{A}$  and  $d_i = \tilde{d}_i$ , we obtain  $A_{ij} = \tilde{A}_{ij} \geq 0$  and the proof is complete.  $\square$

As in Section 2 in the case  $\mathcal{A} = \{0, 1\}$ , it turns out that the roots of the polynomial  $\det(L(G) - xM)$  are simple, provided the mass matrix is well chosen. In order to prepare the proof, we need the following two lemmas.

**Lemma 4** *Suppose  $m = (m_1, \dots, m_n) \in \mathbb{R}^n$  is  $nq$ -independent over a subfield  $K$  of the real numbers and let*

$$\begin{aligned} p_1(x_1, \dots, x_n) &:= \sum_{i=1}^n c_i x_i \\ p_2(x_1, \dots, x_n) &:= \sum_{1 \leq i_1 < i_2 \leq n} c_{i_1 i_2} x_{i_1} x_{i_2} \\ p_3(x_1, \dots, x_n) &:= \sum_{1 \leq i_1 < i_2 < i_3 \leq n} c_{i_1 i_2 i_3} x_{i_1} x_{i_2} x_{i_3} \\ &\vdots \\ p_n(x_1, \dots, x_n) &:= c_{123\dots n} x_1 x_2 \dots x_n \end{aligned}$$

be polynomials with coefficients  $c_{i_1 \dots i_j} \in K \setminus \{0\}$ . Then,  $(p_1(m_1, \dots, m_n), \dots, p_n(m_1, \dots, m_n)) \in \mathbb{R}^n$  is  $q$ -independent over  $K$ .

**Proof**

Let  $F$  be a polynomial in  $n$  variables with coefficients in  $K$  and with maximal degree less than or equal to  $q$ . The maximal degree of  $F$  is defined by

$$\max \deg F := \max_{1 \leq i \leq n} \deg_t F(x_1, \dots, tx_i, \dots, x_n),$$

where  $\deg_t$  is the usual polynomial degree with respect to the variable  $t$ . We assume, that  $F$  is not the zero-polynomial. Now, let us consider the terms in the expression

$$X := F(p_1(m_1, \dots, m_n), \dots, p_n(m_1, \dots, m_n))$$

after expanding all products but before eliminating terms which cancel. We order the  $m$ -monomials in  $X$  according to the lexicographical order relation. E.g.,  $m_1^2 m_2^0 m_3^7 <$

$m_1^2 m_2^1 m_3^1$ . This ordering is compatible with multiplication of the monomials. A lexicographically largest  $m$ -monomial in  $X$  appears while expanding a term

$$p_1^{b_1} p_2^{b_2} \dots p_n^{b_n} = \left( \sum_{i=1}^n c_i x_i \right)^{b_1} \left( \sum_{1 \leq i_1 < i_2 \leq n} c_{i_1 i_2} x_{i_1} x_{i_2} \right)^{b_2} \dots \left( c_{123\dots n} x_1 x_2 \dots x_n \right)^{b_n}$$

(where all exponents  $b_i \leq q$ ) and is, apparently, the monomial

$$m_1^{b_1 + \dots + b_n} m_2^{b_2 + \dots + b_n} \dots m_n^{b_n}$$

(and all exponents here are less than or equal to  $nq$ ). By inspection of this last expression it is clear that the exponent  $(b_1, \dots, b_n)$  is determined by the lexicographically largest  $m$ -monomial in  $X$  which therefore cannot cancel with any other (largest)  $m$ -monomial in  $X$ . Now we assume by contradiction that  $F(p_1(m_1, \dots, m_n), \dots, p_n(m_1, \dots, m_n)) = 0$ . Since all appearing  $m$ -monomials have maximal degree less than or equal to  $nq$ , it follows that all coefficients of the  $m$ -monomials must vanish (because  $(m_1, \dots, m_n)$  is assumed to be  $nq$ -independent over  $K$ ). This contradicts the fact that the coefficient of the lexicographically largest  $m$ -monomial does not vanish.  $\square$

**Lemma 5** *If  $p$  is a polynomial of degree  $n \geq 2$  such that the set of its coefficients is  $(n^2 - 2n + 2)$ -independent over a subfield  $K$  of the real numbers, then  $p$  has only simple roots.*

**Proof**

The polynomial  $p$  has a multiple root if and only if the greatest common divisor of  $p$  and its derivative  $p'$  is non-trivial, i.e., if it is a polynomial of degree strictly larger than zero. The greatest common divisor of two polynomials can be determined by the Euclidean algorithm. Performing the Euclidean algorithm with  $p$  and  $p'$  and computing the polynomial remainders in each step, it is easy to see that the conditions that  $p$  has a multiple root are polynomial equations in the coefficients of  $p$  of degree less than or equal to  $n^2 - 2n + 2$  and with integer coefficients. Since the coefficients of  $p$  are supposed to be  $(n^2 - 2n + 2)$ -independent over  $K \supset \mathbb{Q}$ , the claim follows.  $\square$

**Theorem 5** *Let  $m \in \mathbb{R}_+^n$  be  $n(n^2 - 2n + 2)$ -independent over  $\mathbb{Q}[C]$  and let  $G = (A, M)$  be a connected graph over  $\mathcal{A} = \mathbb{Q}[C]$  of order  $n$ . Then  $\det(L(G) - xM(m))$  has only simple roots.*

**Proof**

Let  $p(x) = a_n x^n + a_{n-1} x^{n-1} + \dots + a_1 x^1$  be as in the proof of Theorem 4. Recall that

$$a_k = \sum_{|I|=k} \det(L_I) \prod_{j \in I} m_j$$

and that  $\det(L_I) \in \mathcal{A} \setminus \{0\}$ , since the graph under consideration is supposed to be connected. Thus, by Lemma 4, we get that  $(a_1, \dots, a_n)$  is  $(n^2 - 2n + 2)$ -independent over  $\mathcal{A}$  and the claim follows from Lemma 5.  $\square$

Combining Theorem 4 and Theorem 5 we obtain

**Theorem 6** *Let  $C \subset \mathbb{R}$  be countable and  $\mathcal{A} = \mathbb{Q}[C]$ . Let  $m \in \mathbb{R}_+^n$  be  $n(n^2 - 2n + 2)$ -independent over  $\mathcal{A}$ , and let  $G = (A, M)$  and  $\tilde{G} = (\tilde{A}, \tilde{M})$  be graphs over  $\mathcal{A}$  of order  $n$  such that  $M$  and  $\tilde{M}$  are permutation similar to  $M(m)$ . Then  $G$  and  $\tilde{G}$  are isomorphic if and only if their Laplace spectra agree, provided that at least one of the graphs  $G$  and  $\tilde{G}$  is connected. In particular, the set of masses  $m \in \mathbb{R}_+^n$  for which reconstruction is not possible is meager and has Lebesgue measure zero.*

**Proof**

Let  $N \subset \mathbb{R}^n$  be the set of all  $m \in \mathbb{R}^n$  which are not  $n(n^2 - 2n + 2)$ -independent over  $\mathbb{Q}[C]$ . Then by Lemma 3 we have that  $N$  is meager and has Lebesgue measure zero. We may assume without loss of generality that  $G$  is connected. By Theorem 5 we have that the roots of  $\det(L(G) - xM(m))$  are all simple. Hence, since the spectra of  $G$  and  $\tilde{G}$  agree, the roots of  $\det(L(\tilde{G}) - x\tilde{M}(m))$  must be simple as well, and the characteristic polynomials of both graphs coincide. Thus, by Theorem 4, the graphs are isomorphic.  $\square$

**Acknowledgment:** We thank the referees for their constructive remarks on the first version of this paper, and we especially thank Walter Gubler for providing the elegant proof of Lemma 4.

## References

- [1] P. Botti, R. Merris: Almost all trees share a complete set of immanental polynomials. *Journal of graph theory* **17** (1993), p. 468ff
- [2] D.M. Cvetković, M. Doob, H. Sachs: *Spectras of graphs*. Academic press, New York (1979)
- [3] B. Eichinger: Configuration statistics of Gaussian molecules. *Macromolecules* **13** (1980), p. 1–11
- [4] F.R.K. Chung: *Spectral graph theory*. CBMS Regional conference series in mathematics; no. 92. American Mathematical Society, Providence, Rhode Island (1997)

- [5] W. Forsman: Graph theory and the statistics of polymer chains. *J. Chemical Phys.* **65** (1976), p. 4111–4115
- [6] R. Grone, R. Merris, V.S. Sunder: The Laplacian spectrum of a graph. *SIAM Journal on matrix analysis and applications* **11** (1990), p. 218
- [7] R. Grone, R. Merris: The Laplacian spectrum of a graph II. *SIAM Journal on discrete mathematics* **7** (1994), p. 221
- [8] I. Gutman: Graph-theoretical formulation of Forsman’s equation. *J. Chem. Phys.* **68** (1978), p. 1321–1322
- [9] L. Halbeisen, N. Hungerbühler: Generation of isospectral graphs. *J. Graph Theory*, to appear
- [10] A.S. Kechris: *Classical descriptive set theory*. Springer, New York (1995)
- [11] D.C. Kurtz: A sufficient condition for all the roots of a polynomial to be real. *Amer. Math. Monthly* **99** (1992), p. 259–263
- [12] B.D. McKay: On the spectral characteristics of trees. *Ars Combinat.* **3** (1977), p. 219–232
- [13] B. Mohar: The Laplacian spectrum of graphs. Preprint series, Dep. of Math. University of Ljubliana, Yugoslavia, **26** (1988), p. 353–384
- [14] J.C. Oxtoby: *Measure and category: a survey of the analogies between topological and measure spaces*. Springer, New York (1980)
- [15] P. Rowlinson: The spectrum of a graph modified by the addition of a vertex. *Publikacije Elektrotehnickog fakulteta* **3** (1992), p. 67ff
- [16] A.J. Schwen: Almost all trees are cospectral. *New directions in the theory of graphs*. Academic press, New York (1973), p. 275–307
- [17] J. Turner: Generalized matrix functions and the graph isomorphism problem. *SIAM J. Appl. Math.* **16** (1968), p. 520–536

Lorenz Halbeisen  
 Dep. of Mathematics  
 U.C. Berkeley  
 Evans Hall 938  
 Berkeley, CA 94720  
 USA  
 halbeis@math.berkeley.edu

Norbert Hungerbühler  
 Dep. of Mathematics  
 University of Alabama at Birmingham  
 452 Campbell Hall  
 1300 University Boulevard  
 Birmingham, AL 35294-1170 USA  
 buhler@mis.mpg.de