

COUNTEREXAMPLES TO F. MOREL'S CONJECTURE ON $\pi_0^{\mathbb{A}^1}$

JOSEPH AYOUB

ABSTRACT. We exhibit counterexamples to F. Morel's conjecture on the \mathbb{A}^1 -invariance of the sheaves of connected components of \mathbb{A}^1 -local spaces.

For a scheme S , we denote by $\mathcal{Spc}(S)$ the ∞ -category $\mathrm{Shv}_{\mathrm{nis}}(\mathrm{Sm}_S)$ of Nisnevich sheaves on smooth S -schemes. An object of $\mathcal{Spc}(S)$ is called an S -space. The Morel–Voevodsky ∞ -category $\mathcal{H}(S)$ is the full sub- ∞ -category of $\mathcal{Spc}(S)$ consisting of \mathbb{A}^1 -local S -spaces. Recall that an S -space \mathcal{X} is \mathbb{A}^1 -local if, for every $U \in \mathrm{Sm}_S$, the map $\mathrm{pr}_1^* : \mathcal{X}(U) \rightarrow \mathcal{X}(U \times \mathbb{A}^1)$ is an equivalence (in the ∞ -category of spaces). The obvious inclusion admits a left adjoint $L_{\mathbb{A}^1} : \mathcal{Spc}(S) \rightarrow \mathcal{H}(S)$.

Notation 1. Let π_0 be the 0-th truncation functor in the ∞ -topos $\mathcal{Spc}(S)$ and, for $i \geq 1$, let π_i be the composition of π_0 with the i -th loop space functor. For an S -space \mathcal{X} , we set $\pi_0^{\mathbb{A}^1}(\mathcal{X}) = \pi_0(L_{\mathbb{A}^1}(\mathcal{X}))$ and, if \mathcal{X} is pointed, we set $\pi_i^{\mathbb{A}^1}(\mathcal{X}) = \pi_i(L_{\mathbb{A}^1}(\mathcal{X}))$.

Now, assume that S is the spectrum of a perfect field k . In his monograph [Mor12], F. Morel proved that the sheaves $\pi_i^{\mathbb{A}^1}(\mathcal{X})$ are \mathbb{A}^1 -invariant in the strongest possible sense for every pointed k -space \mathcal{X} and every integer $i \geq 1$. (See [Mor12, Definition 1.7 & Theorem 1.9] for a precise statement.) The case $i = 0$ was left open and, in [Mor12, Conjecture 1.12], F. Morel expressed the hope that $\pi_0^{\mathbb{A}^1}(\mathcal{X})$ is also \mathbb{A}^1 -invariant for every k -space \mathcal{X} .

We will exhibit counterexamples to F. Morel's conjecture. Interestingly, our counterexamples are based on an old counterexample to a different conjecture of F. Morel, namely his \mathbb{A}^1 -connectivity conjecture over a general base, which we disproved in [Ayo06].

Definition 2. Let X be a smooth k -scheme. We say that X is \mathbb{A}^1 -discrete if, for any extension K/k , every k -morphism $\mathbb{A}_K^1 \rightarrow X$ factors as the structural projection $\mathbb{A}_K^1 \rightarrow \mathrm{Spec}(K)$ followed by a K -point $\mathrm{Spec}(K) \rightarrow X$ of the scheme X .

We have the following well known fact.

Lemma 3. *Let X be a smooth k -scheme. Assume that X is proper and \mathbb{A}^1 -discrete. Then, for a dense open immersion $j : V \rightarrow U$ of smooth k -schemes, composition with j gives a bijection*

$$\mathrm{hom}(U, X) \simeq \mathrm{hom}(V, X).$$

Proof. See [Deb01, Corollary 1.44]. □

We now give a general construction of \mathbb{A}^1 -local k -spaces.

Construction 4. Let X be a smooth k -scheme and let $\mathcal{M} \in \mathcal{H}(X)$ be an \mathbb{A}^1 -local X -space. We denote by $\Phi_X(\mathcal{M})$ the presheaf on Sm_k given informally by

$$U \in (\mathrm{Sm}_k)^{\mathrm{op}} \mapsto \coprod_{s:U \rightarrow X} \Gamma(U; s^* \mathcal{M})$$

Key words and phrases. \mathbb{A}^1 -algebraic topology, F. Morel's conjecture on $\pi_0^{\mathbb{A}^1}$.

where, for every morphism $s : U \rightarrow X$, we write $s^* : \mathcal{H}(X) \rightarrow \mathcal{H}(U)$ for the pullback of \mathbb{A}^1 -local spaces. More precisely, the functor Φ_X is given by the following composition

$$\mathcal{H}(X) \xrightarrow{(\star)} \text{Sect}^{\text{cocart}} \left(\int_{s:U \rightarrow X \in ((\text{Sm}_k)_{/X})^{\text{op}}} \mathcal{H}(U) \Big|_{((\text{Sm}_k)_{/X})^{\text{op}}} \right) \xrightarrow{\Gamma} \text{Psh}((\text{Sm}_k)_{/X}) \xrightarrow{\text{ff}_{X,\sharp}} \text{Psh}(\text{Sm}_k)$$

where $\text{ff}_{X,\sharp}$ is the left Kan extension along the forgetful functor $\text{ff}_X : (\text{Sm}_k)_{/X} \rightarrow \text{Sm}_k$ and (\star) is the obvious equivalence between $\mathcal{H}(X)$ and the ∞ -category of cocartesian sections of the cocartesian fibration classified by $(U \rightarrow X) \mapsto \mathcal{H}(U)$.

Remark 5. Denote by $p : X \rightarrow \text{Spec}(k)$ the structural projection. It can be shown that $L_{\mathbb{A}^1}(\Phi_X(\mathcal{M}))$ is equivalent to $p_{\sharp}(\mathcal{M})$ where $p_{\sharp} : \mathcal{H}(X) \rightarrow \mathcal{H}(k)$ is the left adjoint to the pullback functor p^* . We will not prove this here since we do not need it.

Proposition 6. *Keep the notations as in Construction 4. Assume that X is proper and \mathbb{A}^1 -discrete. Then $\Phi_X(\mathcal{M})$ belongs to $\mathcal{H}(k)$, i.e., it has Nisnevich descent and is \mathbb{A}^1 -invariant.*

Proof. We check the Brown–Gersten property for $\Phi_X(\mathcal{M})$. Clearly, the space $\Phi_X(\mathcal{M})(\emptyset)$ is contractible since $\mathcal{H}(\emptyset)$ is the final category. If $U = U_1 \coprod U_2$, a map $s : U \rightarrow X$ is the union of two maps $s_1 : U_1 \rightarrow X$ and $s_2 : U_2 \rightarrow X$, and we have

$$\Gamma(U; s^* \mathcal{M}) = \Gamma(U_1, s_1^* \mathcal{M}) \times \Gamma(U_2; s_2^* \mathcal{M}).$$

This yields an equivalence $\Phi_X(\mathcal{M})(U) \simeq \Phi_X(\mathcal{M})(U_1) \times \Phi_X(\mathcal{M})(U_2)$. Consider now a Nisnevich square of smooth k -schemes:

$$\begin{array}{ccc} U' & \xrightarrow{j'} & V' \\ \downarrow e' & & \downarrow e \\ U & \xrightarrow{j} & V. \end{array}$$

We need to show that

$$\begin{array}{ccc} \coprod_{s':U' \rightarrow X} \Gamma(U'; s'^* \mathcal{M}) & \longleftarrow & \coprod_{t':V' \rightarrow X} \Gamma(V'; t'^* \mathcal{M}) \\ \uparrow & & \uparrow \\ \coprod_{s:U \rightarrow X} \Gamma(U; s^* \mathcal{M}) & \longleftarrow & \coprod_{t:V \rightarrow X} \Gamma(V; t^* \mathcal{M}) \end{array}$$

is cartesian in the ∞ -category of spaces. Using what we just said, we may assume that V and V' are connected, and that j and j' have dense images. By Lemma 3, we have bijections $\text{hom}(V, X) \simeq \text{hom}(U, X)$ and $\text{hom}(V', X) \simeq \text{hom}(U', X)$. Thus, we may rewrite the above square as follows:

$$\begin{array}{ccc} \coprod_{t':V' \rightarrow X} \Gamma(U'; t'^* \mathcal{M}) & \longleftarrow & \coprod_{t':V' \rightarrow X} \Gamma(V'; t'^* \mathcal{M}) \\ \uparrow & & \uparrow \\ \coprod_{t:V \rightarrow X} \Gamma(U; t^* \mathcal{M}) & \longleftarrow & \coprod_{t:V \rightarrow X} \Gamma(V; t^* \mathcal{M}). \end{array}$$

The obvious map $\text{hom}(V, X) \rightarrow \text{hom}(V', X)$ is injective and the vertical arrows in the above square factor through the summands

$$\coprod_{t:V \rightarrow X} \Gamma(U'; t^* \mathcal{M}) \quad \text{and} \quad \coprod_{t:V \rightarrow X} \Gamma(V'; t^* \mathcal{M})$$

respectively. Thus, we are left to show that the square

$$\begin{array}{ccc} \Gamma(U'; t^* \mathcal{M}) & \longleftarrow & \Gamma(V'; t^* \mathcal{M}) \\ \uparrow & & \uparrow \\ \Gamma(U; t^* \mathcal{M}) & \longleftarrow & \Gamma(V; t^* \mathcal{M}) \end{array}$$

is cartesian for every $t : V \rightarrow X$. This is obvious, since $t^* \mathcal{M}$ belongs to $\mathcal{H}(V)$ by design.

It remains to see that $\Phi_X(\mathcal{M})$ is \mathbb{A}^1 -local. Using that X is \mathbb{A}^1 -discrete, we see that the map $\Phi_X(\mathcal{M})(U) \rightarrow \Phi_X(\mathcal{M})(U \times \mathbb{A}^1)$ is the coproduct over $s : U \rightarrow X$ of the maps

$$\Gamma(U; s^* \mathcal{M}) \rightarrow \Gamma(U \times \mathbb{A}^1; s^* \mathcal{M}).$$

These are equivalences since the $s^* \mathcal{M}$'s belong to $\mathcal{H}(U)$ by design. \square

Next, we describe the sheaves of connected components of the \mathbb{A}^1 -local k -spaces we just constructed.

Proposition 7. *Keep the notations as in Construction 4. Assume that X is proper and \mathbb{A}^1 -discrete. Then the sheaf $\pi_0^{\mathbb{A}^1}(\Phi_X(\mathcal{M}))$ is given by*

$$U \mapsto \coprod_{s:U \rightarrow X} \Gamma(U; \pi_0^{\mathbb{A}^1}(s^* \mathcal{M})).$$

In particular, $\pi_0^{\mathbb{A}^1}(\Phi_X(\mathcal{M}))$ is \mathbb{A}^1 -invariant if and only if $\pi_0^{\mathbb{A}^1}(s^ \mathcal{M})$ is \mathbb{A}^1 -invariant for every morphism $s : U \rightarrow X$. (In particular, a necessary condition is that $\pi_0^{\mathbb{A}^1}(\mathcal{M})$ is \mathbb{A}^1 -invariant.)*

Proof. Since $\Phi_X(\mathcal{M})$ is \mathbb{A}^1 -local, we have $\pi_0^{\mathbb{A}^1}(\Phi_X(\mathcal{M})) = \pi_0(\Phi_X(\mathcal{M}))$. Thus, it is the sheafification of the ordinary presheaf of sets

$$U \mapsto \coprod_{s:U \rightarrow X} \pi_0 \Gamma(U; s^* \mathcal{M}),$$

which we denote by F . Let G be the presheaf described in the statement. We will show that G is a Nisnevich sheaf and that the obvious map $F \rightarrow G$ induces isomorphisms on stalks. This will prove the first statement.

The proof that G is a Nisnevich sheaf is identical to the proof that $\Phi_X(\mathcal{M})$ has Nisnevich descent: we check that G takes a Nisnevich square to a cartesian square of sets, and this boils down to the property that $\pi_0^{\mathbb{A}^1}(t^* \mathcal{M})$ is a Nisnevich sheaf for every $t : V \rightarrow X$, which is true by design. To prove that $F \rightarrow G$ induces an isomorphism on stalks, we fix a henselian essentially smooth k -scheme W . The map $F(W) \rightarrow G(W)$ is then the coproduct, over $r : W \rightarrow X$, of the maps $\pi_0 \Gamma(W; r^* \mathcal{M}) \rightarrow \Gamma(W; \pi_0(r^* \mathcal{M}))$, which are obviously isomorphisms.

For the last statement, using that X is \mathbb{A}^1 -discrete, we see that $\pi_0^{\mathbb{A}^1}(\Phi_X(\mathcal{M}))$ is \mathbb{A}^1 -invariant if and only if, for every $s : U \rightarrow X$, the map

$$\Gamma(U; \pi_0^{\mathbb{A}^1}(s^* \mathcal{M})) \rightarrow \Gamma(U \times \mathbb{A}^1; \pi_0^{\mathbb{A}^1}(s^* \mathcal{M}))$$

is an equivalence. Applying this property for a composite $q \circ s : V \rightarrow X$ with $q : V \rightarrow U$ a smooth morphism, we deduce immediately that the previous condition is equivalent to asking that $\pi_0^{\mathbb{A}^1}(s^* \mathcal{M})$ is \mathbb{A}^1 -invariant for every $s : U \rightarrow X$. \square

It is now clear how to produce counterexamples to F. Morel’s conjecture.

Construction 8. Let X be a smooth, proper and \mathbb{A}^1 -discrete k -scheme. (For example, X can be an abelian variety or a product of curves of genera ≥ 1 .) Let $\mathcal{M} \in \mathcal{H}(X)$ be an \mathbb{A}^1 -local X -space such that $\pi_0^{\mathbb{A}^1}(\mathcal{M})$ is not \mathbb{A}^1 -invariant. Then Proposition 7 ensures that $\pi_0^{\mathbb{A}^1}(\Phi_X(\mathcal{M}))$ is also not \mathbb{A}^1 -invariant. An explicit example of such an \mathcal{M} can be obtained as follows, assuming that X has dimension ≥ 3 . Let $Y \subset X$ be a closed integral surface and $o \in Y(k)$ a rational point admitting a Zariski neighbourhood $N \subset Y$ which is also an étale neighbourhood of the singular point of the projective surface $S \subset \mathbb{P}^3$ defined by the equation $w(x^3 - y^2z) + F(x, y, z) = 0$, where F is a general homogeneous polynomial of degree 4. This is the surface used in [Ayo06] to produce a counterexample to Morel’s connectivity conjecture. In particular, we have a complex of abelian groups $\mathcal{K}_{S,1}^{M,!}$ on Sm_S , concentrated in homological degrees 0 and 1 and sending an irreducible $T \in \text{Sm}_S$ to the two-term Gersten complex

$$k(T)^\times \rightarrow \coprod_{x \in T^{(1)}} \mathbb{Z}.$$

We write also $\mathcal{K}_{S,1}^{M,!}$ for the associated Eilenberg–Mac Lane space which is an object of $\mathcal{H}(S)$. Letting $i : N \rightarrow X$ be the obvious inclusion and $e : N \rightarrow S$ the étale neighbourhood of the singular point of S , we set $\mathcal{M} = i_* e^* \mathcal{K}_{S,1}^{M,!}$. As was shown in [Ayo06], the sheaf $\pi_0^{\mathbb{A}^1}(\mathcal{M})$ restricted to a neighbourhood of o in X is not \mathbb{A}^1 -invariant.

Acknowledgement. This note was written while the author was visiting the Hausdorff Research Institute for Mathematics funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) under Germany’s Excellence Strategy – EXC-2047/1 – 390685813. He thanks the organisers of the trimester program “Spectral Methods in Algebra, Geometry, and Topology” for the invitation and the Hausdorff Research Institute for the excellent working conditions.

REFERENCES

- [Ayo06] Joseph Ayoub. Un contre-exemple à la conjecture de \mathbb{A}^1 -connexité de F. Morel. *C. R. Math. Acad. Sci. Paris*, 342(12):943–948, 2006.
- [Deb01] Olivier Debarre. *Higher-dimensional algebraic geometry*. Universitext. Springer-Verlag, New York, 2001.
- [Mor12] Fabien Morel. \mathbb{A}^1 -algebraic topology over a field, volume 2052 of *Lecture Notes in Mathematics*. Springer, Heidelberg, 2012.

UNIVERSITY OF ZURICH / LAGA - UNIVERSITÉ SORBONNE PARIS NORD
 Email address: joseph.ayoub@math.uzh.ch
 URL: user.math.uzh.ch/ayoub/